
Assessment of farmers' knowledge, attitude and practice of agronomic crop production among smallholder farmers in Imo State, Nigeria

Accessible at: <https://jccr.sccdr.org.ng/index.php/jccr/issue/view/1>

Obinna, Leo O.

Department of Agricultural Extension and Rural Development

Michael Okpara University of Agriculture, Umudike Nigeria

Corresponding Author's Email: obinna.leo@gmail.com

Apu, Uchechi

Department of Agricultural Extension and Rural Development

Michael Okpara University of Agriculture, Umudike Nigeria

Review Process:

Received: 28/11/19

Reviewed: 15/12/19

Accepted: 27/12/19

ABSTRACT

The study assessed the knowledge, Attitude and Practice (KAP) levels of the smallholder farmers in Best Agronomic Practices (BAP) in crop production in Imo State. A sample size of 120 respondents (66.7 % males and 33.3% females) was realized via a multi- stage simple random sampling technique. Data generated via structured questionnaire and interview schedule were analyzed using frequency, percentages, mean and Spearman's correlation coefficient. A mean age of 49 years, a mean household size of 6 persons, mean years of business experience of 16 years, a mean monthly income of ₦30,683.33, and a mean farm size of 0.7 hectares were recorded respectively by the respondents. Seventeen percent of the respondents never had any extension contacts, and 83 % had between once every 2 years and once every month of extension contacts. Results showed high ($X= 55.6$ %) knowledge level, a moderate ($X= 2.3$) attitude level and a low ($X= 1.92$) practice level respectively in BAPs crop production. There was no significant relationship between the knowledge level and practice level in BAP on Crop Production. The study recommends that Government and other stakeholders should improve upon funding and management of extension delivery services in the study area in order for extension to effectively raise the practice level of the respondents in BAPs on crop production as to ensure sustainability of the environment.

Keywords: *Best Agronomic Practices (BAP), Crop Production, Smallholder Farmers.*

INTRODUCTION

The need to improve environmental performance for agriculture, by achieving eco - friendly situations through best agronomic practices in order to ensure sustainability of resource- base has severally been emphasized (Agbarevo, 2016 and Moswetsi, Fanadzo and Ncube, 2017). Agriculture is the most dominant land use and major user of land and water resources in Nigeria (NAERLS), 2017). It creates employment for about 70% of rural dwellers who engage in it and its' related activities, it produces primary materials for

industries, provides food for animals and humans, and also the second foreign exchange earner after oil for the country Nigeria (Ekong, 2010). Therefore, for agriculture to remain viable within the conservation needs of the farm, surroundings and management techniques must be able to preserve and restore the critical resource – base (soil and water) as to ensure sustainability

Similarly, Ekong (2010) noted that agriculture in the South Eastern Zone of Nigeria is mostly rain – fed, small- scale and fallow extractive in nature therefore, that any change in environment is bound to impact on productivity and other socio- economics activities of the farmers in the zone. On that note, Obinna (2015) observed that farmers in the South Eastern Zone of Nigeria are confronted with the problems of agricultural land scarcity caused by high population densities, diversion of agricultural lands to other uses, coupled with the soil that is structurally fragile and burdened with unfavourable land tenure situations, and over dependency of agricultural activities on the vagaries of the weather, aggravated by inconsistency in the onset of the rains and their distributions during the rainy season among other limiting factors have compelled the farmers in the zone to farm on marginal lands in an unsustainable manner thereby degrading the soil and polluting the water- resources.

In addition, Asiabaka (2012) noted that in Imo State due to pressure of population growth on lands and other factors that the traditional methods of maintaining soil fertility by organic fertilizers and use of trees in many parts of the State have disappeared partly or completely. As such soil degradation and erosion take place in a large scale, but also that traditional knowledge and skills as well as indigenous genetic resources practices are being eroded away. Several studies have equally, shown that in recent times, poverty level and hunger are in the increase in Nigeria and Imo state inclusive. This has been attributed to poor farm returns, high cost of labour, lack of funding for ADPs to carry out their extension work effectively, high inflation rates, and high cost for food items, which have made Imo state to be food insecure (Okoroh, Olaolu and Igbokwe, 2016).

It is in the light of the above that the study sought to assess the gap in knowledge, Attitude and Practice (KAP) levels of the respondents in Best Agronomic Practices (BAPs) in crop production in the study area. The following objectives guided the study:

- (i) examine the socio- economic characteristics of the smallholder farmers;
- (ii) ascertain the knowledge level of the respondents in BAPs in crop production;
- (iii) ascertain the attitude level of the respondents in BAPs in crop production, and
- (iv) ascertain the practice level of the respondents in BAPs in crop production.

It was hypothesized that there was no significant relationship between the knowledge level of the respondents and their practice level in BAPs in crop production in the study area.

METHODOLOGY

The study was conducted in Imo State, Nigeria. Imo State is located in the South- Eastern Ecological Zone of Nigeria. It lies between Longitude 6° 50' and 7° 25' East of the Greenwich Meridian and Latitude 4° 45' and 7° 15' North of the Equator. A multi- stage and simple random method was employed alongside purposive sampling technique in selecting a sample size of 120 respondents. Stage 1; the 3 Agricultural Zones of the state were purposively selected. Stage 2, made use of simple random method, to select 2 Agricultural Blocks from each of the 3 Agricultural Zones to give 6 Agricultural Blocks. Stage 3, also made use of simple random method, to select 2 Agricultural Cycles from each of the 6 Agricultural blocks to give 12 cycles. Stage 4, equally, used simple random method with

the assistance of Extension Agents, to select 10 respondents from each of the 12 cycles to give a total of 120 respondents that were used for the study. Primary data were generated via the use of questionnaire and interview schedule and were analyzed using frequency counts, percentages, means, pooled means and Spearman's Coefficient of Correlation.

The gap in knowledge, Attitude and Practice (KAP) level of the respondents in BAPs in crop production in the study area stressed the need to provide strategic, critical, and "quality" information, which included non-technological information as the reasons why low practice level of best agronomic practices in crop production which might be related to socio-psychological, socio-cultural, and socio-economic factors. It was based on this that the respondents' KAP levels in BAPs were weighted and scored as follows: The knowledge level of the respondents was measured through a 16 item statements, where the respondents were asked to indicate if they were aware scored one point and if not aware scored zero point in all the 16 item statements in BAPs in crop production in the study area. A percentage score of all the positive responses of the respondents on the 16 item statements was computed based on the number scored divided by 16 and multiplied by 100 to give percentage score for each of the BAPs.

A bench score of 50% was established. Any % score $\geq 50\%$ was regarded as a knowledgeable practice while any % score $< 50\%$ was regarded as not knowledgeable practice. The Knowledge level of the respondent was categorized as follows: From 0% - 49% = low knowledge level, and 50% - 100% = high knowledge level. Attitude of the respondents in BAPs was measured on 16 item statements as well in BAPs through a 4 point Likert type scale of strongly agree, weighted 4 points, agree weighted 3 points, disagree weighted 2 points and strongly disagree weighted 1 point respectively. A bench mark of 2.5 was established and any of the BAPs that had any mean score ≥ 2.5 was regarded as positive attitude while on the other hand any mean score < 2.5 was regarded as negative attitude. The attitudinal level of the respondents in BAPs in crop production was categorized as follows: 0.0 - 0.99 = very low attitude level. 1.00 - 1.99 = low attitude level 2.00 - 2.99 = moderate attitude level and 3.00 - 4.00 = high attitude level. The practice level was measured through a 4 point Likert type scale of always weighted 4 points, often weighted 3 points, sometime weighted 2 points and never weighted 1 point respectively. A mean of 2.5 was established in that any mean score ≥ 2.5 was regarded as practiced while any mean score < 2.5 was regarded as not practiced. The level of practice of the respondents was categorized as follows:

- 0.00 - 0.99 very low level of practice
- 1.00 - 1.99 low practice level
- 2.00 - 2.99 moderate practice level
- 3.00- 4.00 high practice level

RESULTS AND DISCUSSION

Socioeconomic Characteristics.

The Socio-Economic Characteristics is as presented in Table 1. Table 1 shows that the mean age of the respondents was 49 years, 66.7% were males and 33.3% females and 15% were single, 75% married, 4.2% divorced / separated and 5.8% widows respectively. This implied that the respondents were mature, responsible and at the peak of their socio-economic activities (Ekong, 2010). About 91.7% of the respondents were literates. This implied that a very high proportion (91.7%) of the respondents could read and write. Table 1 equally shows that 41.7% of the respondents were farmers, 48.3% traders, 1.7% civil

servants and 8.3 % other professions. The higher percent (58.3 %) of the respondents that were involved in non- farming activities confirmed that there is problem of farm- land scarcity and poor farm returns confronting them in the study area which have forced them to go into non- farming activities in order to make ends meet. A mean household size of 6 persons was recorded. This is very typical of the rural households in the Southern part of Nigeria as postulated by Ekong (2010). A mean monthly income of ₦ 30,683.33 was equally recorded. This could be due to the fact that a reasonable proportion (58.3%) of the respondents were involved in non- farming activities which gave higher monthly returns. Table 1 further recorded a mean farm size of 0.7 hectares. This could be due to scarcity of farm- lands because of population pressure and diversion of agricultural land to other uses. About 16 years of mean years of business experience was recorded and 25% of the respondents were into mixed-farming, and 50 %, 6.3 %, 8.3% and 3.3% were into crop- production, livestock production, bee-keeping, fisheries, snailery respectively. The implication was that the respondents were into all these types of farming activities due to scarcity of farm land that is characteristic of the Zone and also to diversify as safety net measures since returns from farming are poor and inconsistent (Obinna, 2015). About 12.5% of the respondents practiced shifting cultivation, 16.7%, 37.5 %, 12.5 % and 20.8% were into fallow / crop rotation, continuous cultivation, alley – farming, and agro- forestry respectively. Respondents in the study area practiced all these types of cultivations as to find solution to the problem of soil degradation that is very common. Table 1 finally shows that 16.7 % of the respondents had no extension contacts, and about 83.3% had between once every 2 years and once every month of extension contacts. This might be due to poor funding of Extension by the ADP in the study area which has made the extension not to carry out its function effectively.

Table 1: Distribution of the Respondents According to Socio- economic Characteristics (n= 120)

| Variables | Frequency | Percentage | Cumulative Frequency | Mean |
|-------------------------------|-----------|------------|----------------------|----------|
| Age in years | | | | |
| ≤ 25 | 08 | 6.67 | 6.67 | |
| 26 - 36 | 15 | 12.50 | 19.17 | |
| 37 - 47 | 28 | 23.33 | 42.50 | 49 years |
| 48 - 58 | 32 | 26.67 | 69.17 | |
| 59 - 69 | 25 | 20.83 | 90.00 | |
| 70 & above | 12 | 10.00 | 100.00 | |
| Sex | | | | |
| Male | 80 | 66.67 | 66.67 | |
| Female | 40 | 33.33 | 100.00 | |
| Marital Status | | | | |
| Single | 18 | 15.00 | 15.00 | |
| Married | 90 | 75.00 | 90.00 | |
| Separated/ Divorced | 5 | 4.17 | 94.17 | |
| Widow | 7 | 5.83 | 100.00 | |
| Educational Attainment | | | | |
| No formal education | 10 | 8.33 | 8.33 | |
| Primary school completed | 60 | 50.00 | 58.33 | |
| Sec School completed | 40 | 33.33 | 91.66 | |
| Higher education completed | 10 | 8.33 | 99.99 | |

Table 1 continued

| Variables | Frequency | Percentage | Cumulative Frequency | Mean |
|--|-----------|------------|----------------------|---------------|
| Occupation | | | | |
| Farming | 50 | 41.67 | 41.67 | |
| Trading | 58 | 48.33 | 90.00 | |
| Civil service | 02 | 1.67 | 91.67 | |
| Other profession | 10 | 8.33 | 100.00 | |
| Household Size | | | | |
| ≤ 4 | 30 | 25.00 | 25.00 | |
| 5 - 7 | 60 | 50.00 | 75.00 | 6 persons |
| 8 & above | 30 | 25.00 | 100.00 | |
| Monthly income (₦) | | | | |
| ≤ 20,000.00 | 30 | 25.00 | 25.00 | |
| 21,000.00 - 31,000.00 | 40 | 33.33 | 58.33 | |
| 32,000.00 - 42,000.00 | 30 | 25.00 | 83.33 | |
| 43,000.00 - 53,000.00 | 10 | 8.33 | 91.66 | ₦30,683.33 |
| 54,000.00 - 64,000.00 | 08 | 6.67 | 98.33 | |
| 65,000.00 & above | 02 | 1.67 | 100.00 | |
| Farm size in hectares | | | | |
| ≤ 0.1 | 30 | 25.00 | 25.00 | |
| 0.2 - 0.7 | 40 | 33.33 | 58.33 | |
| 0.8 - 1.3 | 30 | 25.00 | 83.33 | 0.79 hectares |
| 1.4 & above | 20 | 16.67 | 100.00 | |
| Farming Experience in yrs | | | | |
| ≤ 10 | 10 | 8.33 | 8.33 | |
| 11 - 21 | 30 | 25.00 | 33.33 | |
| 22 - 32 | 40 | 33.33 | 66.66 | 16.17 years |
| 34 & above | 40 | 33.33 | 99.99 | |
| Type of Farming | | | | |
| Mixed Farming | 30 | 25.00 | 25.00 | |
| Crop Production | 60 | 50.00 | 75.00 | |
| Livestock Production | 08 | 6.33 | 81.33 | |
| Fisheries | 10 | 8.33 | 89.66 | |
| Bee- keeping | 08 | 6.33 | 95.99 | |
| Snailery | 04 | 3.33 | 100.00 | |
| Farming System | | | | |
| Shifting cultivation | 15 | 12.50 | 12.50 | |
| Fallow/ crop rotation | 20 | 16.67 | 29.17 | |
| Continuous cultivation | 45 | 37.50 | 66.67 | |
| Alley farming | 15 | 12.50 | 79.17 | |
| Agro- forestry | 25 | 20.83 | 100.00 | |
| Frequency of Extension contacts | | | | |
| None | 20 | 16.67 | 16.67 | |
| Once every 2 years | 20 | 16.67 | 33.34 | |
| Once every year | 15 | 12.50 | 45.84 | |
| Once every 6 months | 25 | 20.83 | 66.67 | |
| Once every 3 months | 20 | 16.67 | 83.34 | |
| Once every month | 20 | 16.67 | 100.0 | |

Source Field Survey 2019

Determination of Knowledge Level of the Respondents in BAPs in Crop Production

The Knowledge Level in BAPs in Crop Production is as presented in Table 2. Table 2 shows that the respondents had high ($\bar{X} \geq 50\%$) knowledge level on 12 item statements of BAPs in crop production out of 16 item statements. The ones they indicated low ($\bar{X} = 0 - 49\%$) knowledge level included knowledge on correct plant geometry ($\bar{X}=41.66\%$), recommended tillage operations for each crop ($\bar{X}= 45.83\%$), mulching technique against water loss ($\bar{X} = 45.83\%$) and optimum cropping patterns ($\bar{X} = 45.83\%$) respectively. It implied that these were highly technical innovations that normally come from the scientists' research stations and transmitted to the farmers by the Extension Agents. Therefore, in conclusion, the respondents in the study area had a high ($\bar{X} \geq 50\%$) knowledge level in BAPs on crop production since the grand mean ($\bar{X} = 55.63\%$).

Table 2: Distribution of the Respondents According to Knowledge level in BAPs Crop Production (n= 120)

| S/No | Best- Agronomic Practices | Frequency | Percentage scores | Knowledge Level |
|------|---|-----------|-------------------|-----------------|
| 01 | Do you know about improved cultivar | 80 | 66.67 | High |
| 02 | Do you know about recommended planting dates for each crop | 70 | 58.33 | High |
| 03 | Do you know about the correct planting population for each crop | 70 | 58.33 | High |
| 04 | Do you know about correct plant geometry as recommended | 50 | 41.66 | Low |
| 05 | Do you know about the recommended tillage operations for each crop | 55 | 45.83 | Low |
| 06 | Do you know that you should incorporate plant residues in your farm | 60 | 50.00 | High |
| 07 | Do you know about mulching of your crops against the loss of water | 55 | 45.83 | Low |
| 08 | Do you know about optimum water and fertilizer management | 60 | 50.00 | High |
| 09 | Do you know about optimum cropping pattern for each crop | 55 | 45.83 | Low |
| 10 | Do you know about farm yard, cow dung, poultry droppings and compost manure | 75 | 62.50 | High |
| 11 | Do you know about integrated pest management system (IPM) | 60 | 50.00 | High |
| 12 | Do you know about cover cropping system of farming | 70 | 58.33 | High |
| 13 | Do you know about alley- farming | 70 | 58.33 | High |
| 14 | Do you know about agro - forestry system of farming | 80 | 66.67 | High |
| 15 | Do you know about liming to improve the texture and structure of the soil | 70 | 58.33 | High |
| 16 | Do you know about irrigation | 90 | 75.00 | High |

Source: Field Survey 2019

Determination of the Attitude Level of the Respondents in BAPs in Crop Production in the Study Area.

The Attitude Level in BAPs in Crop Production is as presented in Table 3 below. Table 3 shows that the respondents had low ($\bar{X}= 1.00 - 1.99$) attitude level in 6 item statements in BAPs in crop production. They include: following the correct planting dates increases

yield ($\bar{X} = 1.83$), following the correct planting density (population) increase yield ($\bar{X}=1.75$), following the recommended tillage operations increase yield ($\bar{X}= 1.63$), following the optimum cropping pattern increases yield ($\bar{X}= 1.79$), use of organic manure improves soil structure and increases yield.

$\bar{X}= 1.83$), and integrated pest management (IPM) is the best way to sustain the environment ($X=1.96$). The respondents equally, indicated moderate ($\bar{X}= 2.00 - 2.99$) attitude level in 8 items statements out of 16 item statements. They include: planting improved cultivar is the best way to increase yield ($\bar{X}=2.66$), following the correct plant geometry increases yield ($\bar{X}= 2.00$), incorporating plant residues increases soil fertility ($\bar{X}= 2.50$), mulching your crops protect your crop against water loss ($\bar{X}= 2.67$), optimum fertilizer and water management is the best in farming ($\bar{X}=2.92$), cover- cropping system improves soil fertility ($\bar{X}= 2.29$), alley- farming protects and enriches the soil ($\bar{X}=2.71$), and agro- forestry system is the most sustainable way of farming in fragile soil ($\bar{X}=2.63$). Table 3 equally shows that Farm yard residue, cow dung & poultry droppings improve the soil texture and structure and irrigation allows you to farm during the dry season respectively recorded high ($\bar{X}= 3.08$) attitude level in BAPs in crop production. Table 3 finally shows that the grand mean ($\bar{X}= 2.3$) attitude level which implies that holistically the respondents in the study area had a moderate ($\bar{X}= 2.00 - 2.99$) attitude level in BAPs in crop production in the study area.

Table 3: Distribution of the Respondents According to their Attitude Level on Best – Agronomic Practices in Crop Production (n= 120)

| S/No | Best- Agronomic Practices | Strongly agree | Agree | Disagree | Strongly Disagree | Mean | Attitudinal Level |
|------|--|----------------|-------|----------|-------------------|------|-------------------|
| 01 | Planting improved cultivar is the best way to increase yield | 30 | 40 | 30 | 20 | 2.66 | Moderate |
| 02 | Following the recommended planting dates increases yield | - | 20 | 60 | 40 | 1.83 | Low |
| 03 | Following the correct plant population increases yield | - | 30 | 30 | 60 | 1.75 | Low |
| 04 | Correct plant geometry increases yield | - | 40 | 40 | 40 | 2.00 | Moderate |
| 05 | Following recommended tillage operations increases yield | - | 35 | 35 | 50 | 1.63 | Low |
| 06 | Incorporating plant residues increases soil fertility | 30 | 30 | 30 | 30 | 2.50 | Moderate |
| 07 | Mulching of your crops help against water loss | 30 | 40 | 30 | 20 | 2.67 | Moderate |
| 08 | Optimum water and fertilizer management is the best in farming | 40 | 40 | 30 | 10 | 2.92 | Moderate |
| 09 | Optimum cropping pattern increases yield | - | 25 | 45 | 50 | 1.79 | Low |
| 10 | The use of organic manure and compost improves the soil & increases yield | 10 | 20 | 30 | 60 | 1.83 | Low |
| 11 | Integrated pest management system(IPM) is the best way to sustain the environment | 10 | 20 | 45 | 45 | 1.96 | Low |
| 12 | Cover cropping system improves the soil fertility | 20 | 30 | 35 | 35 | 2.29 | Moderate |
| 13 | Alley- farming protects and enriches the soil | 30 | 35 | 35 | 30 | 2.71 | Moderate |
| 14 | Agro – forestry system is the most sustainable way of farming on fragile soil | 35 | 35 | 20 | 30 | 2.63 | Moderate |
| 15 | Farm yard residue, cow dung & poultry droppings improve the soil texture and structure | 50 | 40 | 20 | 10 | 3.08 | High |
| 16 | Irrigation allows you to farm during the dry season | 55 | 35 | 15 | 15 | 3.08 | High |

Source: Field Survey 2019

Determination of the Practice Level of the Respondents in BAPs in Crop Production

The Practice Level in BAPs in Crop Production is as presented in Table 4 below. Table 4 shows that the respondents had low (1.00 - 1.99) practice level in 9 item statements in BAPs out of 16 item statements in BAPs in crop production. Table 4 further shows that the respondents had moderate (\bar{X} = 2.00 - 2.99) practice level in in 7 item statements out of 16 item statements in BAPs in crop production. Table 4 equally shows the grand mean (\bar{X} = 1.92) which implies that the respondents generally had a low (\bar{X} = 1.00 - 1.99) practice level in BAPs in crop production in the study area. This might be linked to the low extension contacts among the respondents in the study area.

Table 4: Distribution of the Respondents According to their Practice Level in BAPs in Crop Production in the Study Area (n = 120)

| S/No | Best- Agronomic Practices in Crop Production | Always | Often | Sometimes | Never | Mean | Practice Levels |
|------|--|--------|-------|-----------|-------|------|-----------------|
| 01 | How often do you plant improved cultivar? | 5 | 15 | 35 | 65 | 1.66 | Low |
| 02 | How often do you plant according to the recommended planting dates? | - | 20 | 40 | 60 | 1.66 | Low |
| 03 | How often do you plant according to the correct plant population (plant density) ? | - | 25 | 35 | 60 | 1.71 | Low |
| 04 | How often do you plant according to the correct plant geometry? | 5 | 20 | 30 | 65 | 1.71 | Low |
| 05 | Do you always follow the correct tillage operations for your crops? | 10 | 20 | 30 | 60 | 1.66 | Low |
| 06 | Do you always incorporating plant residues into the soil? | 15 | 20 | 35 | 50 | 2.00 | Moderate |
| 07 | Do you always carry out mulching of your crops against water loss? | 20 | 30 | 40 | 30 | 2.33 | Moderate |
| 08 | Do you always practice optimum water and fertilizer management in your farming? | 20 | 25 | 35 | 40 | 2.21 | Moderate |
| 09 | How often do you practice optimum cropping pattern? | 10 | 20 | 30 | 60 | 1.83 | Low |
| 10 | Do you always use organic manure and compost in your farm? | 20 | 30 | 30 | 40 | 2.25 | Moderate |
| 11 | Do you always practice integrated pest management system (IPM) in your farming operations? | - | 15 | 20 | 85 | 1.58 | Low |
| 12 | How often do you carry out cover cropping in your farming system? | 5 | 10 | 25 | 80 | 1.54 | Low |
| 13 | How often do you practice alley-farming? | 10 | 20 | 30 | 60 | 1.83 | Low |
| | How often do you practice agro – forestry? | 20 | 30 | 40 | 30 | 2.33 | Moderate |
| 15 | How often do you use farm yard residue, cow dung & poultry droppings improve the soil texture and structure? | 25 | 30 | 35 | 30 | 2.42 | Moderate |
| 16 | Do you always practice irrigation for dry season farming? | 15 | 25 | 35 | 45 | 2.08 | Moderate |

Source: Field Survey 2019

Hypothesis Testing

There was no significant relationship between the knowledge level of the respondents and their practice level in BAPs in crop production in the study area since $\Gamma_s = 0.955$. Therefore, H_0 is accepted.

CONCLUSION

The results showed that the respondents had a high ($\bar{X} = 55.6\%$) knowledge level in BAPs and a moderate ($\bar{X} = 2.3$) attitude level and a low ($\bar{X} = 1.92$) practice level in BAPs in crop production. The results equally showed that there was no significant relationship between the knowledge level in BAPs and practice level of the respondents in crop production in the study area. The study therefore concluded that the respondents had a low ($X = 1.92$) practice level in BAPs in crop production in the study area due to poor extension contacts to them. The study therefore recommended that Government and other stakeholders in extension delivery services should improve upon funding and management of extension delivery services in the study area in order for extension to effectively raise the practice level of the respondents in BAPs on crop production as to ensure sustainability of the environment.

REFERENCES

- Agbarevo, M. N. B (2016) Farmers' Adoption of Sustainable Agronomic Practices in Cross- River State, Nigeria, Proceedings, 21st Annual National Conference of the Agricultural Extension Society of Nigeria 2016 pp 269 – 275.
- Ahaneku, I. E. (2010). Conservation of Soil and Water- resources for Combating Food crisis in Nigeria. Scientific Research and Essays 5(6), pp. 507 – 573, *Academic Journals*.
- Asiabaka, C.C. (2012). Promoting Indigenous Knowledge (IK) for the Attainment of the Millennium Development Goals (MDGs). A key note paper presented to the Nigerian Society for Indigenous Knowledge and Development during the 6th Annual Conference on 5th – 7th June 2012 at MOUUA Abia State, Nigeria.
- Ekong, E. E. (2010) Rural Sociology (3rd Ed) Dove Educational Publishers Uyo Nigeria pp 1-425
- Moswetsi, G; Fanadzo, M and Ncube, B (2019). Cropping Systems and Agronomic Management Practices in Smallholder Farms in South Africa Constraints, Challenges and Opportunities. *Journal of Agronomy*, 16 pp 51 – 64 Retrieved on the 20/02 / 2019
- National Agricultural Extension Research Liaison Services (NAERLS) (2017). N-Power Agro-Training Manual for Extension Advisors, 1 – 354. Editors; M. K, Othman, A. O, Lawal and A. O, Iyiola- Tunji. Federal Ministry of Agriculture and Rural Development, Abuja.
- National Population Commission (NPC) (2007). A Blueprint 2006 National Census Abuja, Nigeria
- Obinna, Leo. O. (2015): Effect of Cassava- Value Chain on Income of Small- Scale Farmers in Abia State, Nigeria. *Discourse Journal of Agriculture and Food Science* Vol.3 (9): 128 – 134 September, 2015.
- Okoroh, J. P.; Olaolu, M. O and Igbokwe, E. M (2016). Climate Change Mitigation and Adaptation Strategies Used by Farmers in Imo State, *Nigeria Journal of Agricultural Extension* Vol. 20 (2) pp 129 - 141 <http://journal.aessonigeria.org>
- Qzor, N.; Madukwe, M. C.; Enete, A. A.; Amaechina, E.C.; Onokala, P.; Eboh, E. C.; Ujah, O. and Garforth, C. J (2010). Barriers to Climate Change Adaptation Among Farming Households of Southern Nigeria. *Journal of Agricultural Extension* Vol. 14(1) pp 131 -143 June, 2010
- Sinkaiye, T (2011). Participatory Methodologies and Approaches in Agricultural Extension In Agricultural Extension in Nigeria AESON 2nd Edition pp 294 - 309 AESON Nigeria.